

PATENT APPLICATION

OF

WILLIAM J. ROBERTS

FOR

**“BIOAVAILABLE PRODRUGS OF ANDROGENIC STEROIDS
AND RELATED METHOD”**

CERTIFICATE OF EXPRESS MAIL

Express Mail Mailing Label No. EV054088650US

Date of Deposit: January 16, 2002

I hereby certify that this paper or fee is being deposited with the United States Postal Service “Express Mail Post Office to Addressee” service under 37 CFR 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner for Patents, U.S. Patent and Trademark Office, P.O. Box 2327, Arlington, VA 22202-0327.

Anne K Hoole

Attorney Docket No. 1400-003

(Atty Docket 1400-003)

5 BACKGROUND OF THE INVENTION

Field of the Invention

10 Description of the Related Art

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846

[0003] A known approach to addressing such steroid concentration deficiencies is to introduce the deficient steroid into the body. In U.S. Patent No. 5,578,588, for example, methods are disclosed for delivering testosterone

supplements including peroral administration or intramuscular injection of testosterone.

[0004] This approach, however, suffers from a number of disadvantages.

Introduction of such steroid supplements has been associated with undesirable effects such as poor control over blood concentrations and loss of the steroid due to a “first pass effect,” wherein the steroid is metabolized by the liver prior to reaching general circulation. These losses dramatically reduce the available steroid and, consequently, much higher doses of the steroid supplement generally must be administered to achieve the desired effects. The higher doses sometimes result in an undesirable and unpredictable rise in overall steroid concentration, which, for example, in the case of testosterone, can result in physiological and psychological problems.

[0005] Another approach to addressing steroid concentration deficiencies is to introduce a prodrug of the steroid into the body. An example of a prodrug is a prohormone. A prohormone is a compound that itself has no anabolic activity but, when administered in the body, is metabolized or converted into a natural or desired hormone. Such prohormones become substrates for in vivo bioconversion into the parent compounds, i.e., the corresponding natural or desired hormones.

U.S. Patent No. 5,053,403, for example, discloses that specific prohormones including androstenedione, progesterone, and 17α - β derivatives or analogues can be administered to humans for the purpose of increasing blood concentration of

1006340307

testosterone, with fewer undesirable effects. Long term use of these prohormones, however, is also associated with side effects, such as gynecomastia. A pernasal dose of 3.5 mg to 15 mg of these prohormones is reported to increase the blood concentration of testosterone by 34% to 97%. Similarly, U.S. Patent No. 5,880,117
5 discloses the use of 4-androstenediol as a peroral testosterone supplement. Androstenedione is a direct precursor of testosterone and estrogen in target tissues having appropriate receptors and enzymes. According to U.S. Patent No. 6,117,429, androstenediols are precursors for testosterone after oral administration in adults. 19-norandrostenedione is a precursor for 19-nortestosterone, which has a similar anabolic activity in comparison to testosterone.

[0006] The general approach of using prodrugs or prohormones to achieve supplementation of androgenic steroid concentrations in vivo also has been limited, however, in that the effectiveness of such compounds has tended to be low. In some instances, their conversion into the desired steroid is limited, for example, because
15 they are removed from the system through the first pass effect. They also can be converted into undesirable products, for example, as in the case wherein 4-androstenedione is converted into estrogen. Even where the desired bioconversion occurs, the rate of conversion can be sufficiently low that undesirably large quantities of the prodrug must be taken to achieve desired results. This itself can
20 have undesirable side effects.

SUMMARY OF THE INVENTION

[0011] To achieve the foregoing objects, and in accordance with the purposes of the invention as embodied and broadly described in this document, a compound is provided for increasing the concentration of a parent androgen in a subject in vivo, wherein the parent androgen has a skeletal structure including a 1 position and a 17 position and the parent androgen further has a 17 β -hydroxy group comprising a 17 β -hydroxy oxygen appended to the 17 position and a 17 β -hydroxy hydrogen appended to the 17 β -hydroxy oxygen.

[0012] The compound according to this aspect of the invention comprises a substrate having the skeletal structure of the parent androgen comprising a 1 position and a 17 position corresponding to the 1 and 17 positions respectively of the parent androgen. The substrate further comprises a carbon-carbon double bond at the 1 position. The compound also comprises a promoiety appended to the 17 position of the substrate as a substitute for the 17 β -hydroxy hydrogen of the parent androgen. The promoiety comprises an alkylcarbonate ester.

[0013] The substrate has the skeletal structure of the parent androgen, preferably wherein the parent androgen is selected from the group consisting of 5 α -androst-1-ene-3 α ,17 β -diol, 5 α -androst-1-ene-3 β ,17 β -diol, and mixtures thereof. The substrate also may have the skeletal structure of the parent androgen wherein the parent androgen is selected from the group consisting of 5 α -estr-1-ene-3 α ,17 β -diol, 5 α -estr-1-ene-3 β ,17 β -diol, and mixtures thereof. In addition, the parent androgen

5

to which the substrate corresponds may comprise 17 β -hydroxy-5 α -androst-1-ene-3-one. The substrate still further may have the skeletal structure of the parent androgen wherein the parent androgen is selected from the group consisting of 17 β -hydroxyandrost-1,4-diene-3-one, 17 β -hydroxyestr-1,4-diene-3-one, 17 β -hydroxy-5 α -estr-1-ene-3-one, and mixtures thereof.

15

[0014] The alkylcarbonate ester optionally but preferably has an alkyl chain length of less than 12, and more preferably of less than 11. The alkylcarbonate ester may be selected from the group consisting of methyl carbonate, ethyl carbonate, propyl carbonate, isopropyl carbonate, butyl carbonate, isobutyl carbonate, t-butyl carbonate, valeryl carbonate, hexyl carbonate, heptyl carbonate, octyl carbonate, nonyl carbonate, decyl carbonate, undecyl carbonate, dodecyl carbonate, cyclopentyl carbonate, cyclopentylmethyl carbonate, cyclopentylpropyl carbonate, cyclohexylmethyl carbonate, cyclohexylpropyl carbonate, and mixtures thereof. The compound itself according to this aspect of the invention may assume a number of specific forms. It may, for example, comprise 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -alkylcarbonate, 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate, 17 β -hydroxyandrost-1,4-diene-3-one 17 β -alkylcarbonate, or 17 β -hydroxyandrost-1,4-diene-3-one 17 β -ethylcarbonate. It also may take the form of mixtures or combinations of these. Incidentally, compounds identified or referred to herein that identify a bond or functional group at the 3 position, e.g., those identified in this paragraph, may comprise a 3 α or a 3 β unless indicated otherwise.

5 [0015] The compound also may take the form of estranes and derivatives, such as ketone derivatives. It may, for example, comprise 17 β -hydroxy-5 α -estr-1-ene-3-one 17 β -alkylcarbonate, 17 β -hydroxyestr-1,4-diene-3-one 17 β -alkylcarbonate, and like forms. The ethyl carbonates are preferred forms of the alkylcarbonate promoiety in such molecules, e.g., 17 β -hydroxy-5 α -estr-1-ene-3-one 17 β -ethylcarbonate.

10 [0016] Compounds according to this aspect of the invention also may comprise diols. Examples would include 5 α -androst-1-ene-3-ol 17 β -alkylcarbonate, 5 α -androst-1-ene-3,17 β -diol 17 β -alkylcarbonate (e.g., 5 α -androst-1-ene-3,17 β -diol 17 β -ethylcarbonate), 5 α -androst-1-ene-3,17 β -diol-3,17 β -di(alkylcarbonate) (e.g., 5 α -androst-1-ene-3,17 β -diol-3,17 β -di(ethylcarbonate)), and similar forms.

15 [0017] In some forms, particularly those intended for peroral administration, it is preferable albeit optional for the compound to include a carrier. The carrier may be a solid, a liquid, a semi-solid, or other suitable form.

20 [0018] In accordance with another aspect of the invention, a method is provided for increasing the concentration of a parent androgen in a subject in vivo, wherein the parent androgen has a skeletal structure including a 1 position and a 17 position and the parent androgen further has a 17 β -hydroxy group comprising a 17 β -hydroxy oxygen appended to the 17 position and a 17 β -hydroxy hydrogen appended to the 17 β -hydroxy oxygen.

[0019] The method according to this aspect of the invention comprises administering to the subject a compound comprising a substrate and a promoiety. The substrate has the skeletal structure of the parent androgen comprising a 1 position and a 17 position corresponding to the 1 and 17 positions respectively of the parent androgen, and the substrate comprises a carbon-carbon double bond at the 1 position. The promoiety is appended to the 17 position of the substrate as a substitute for the 17 β -hydroxy hydrogen of the parent androgen, and comprises an alkylcarbonate ester. The method further comprises converting the compound in vivo into the parent androgen. The subject may be a human being, in which case the in vivo conversion comprises converting the compound into the parent androgen in vivo within the human being.

[0020] In accordance with this aspect of the invention, the skeletal structure of the parent androgen to which the substrate corresponds may be selected from the group consisting of 5 α -androst-1-ene-3 α ,17 β -diol, 5 α -androst-1-ene-3 β ,17 β -diol, and mixtures thereof. It also may be selected from the group consisting of 5 α -estr-1-ene-3 α ,17 β -diol, 5 α -estr-1-ene-3 β ,17 β -diol, and mixtures thereof. The skeletal structure of the parent androgen to which the substrate corresponds also may comprise 17 β -hydroxy-5 α -androst-1-ene-3-one. It also may be selected from the group consisting of 17 β -hydroxyandrost-1,4-diene-3-one, 17 β -hydroxyestr-1,4-diene-3-one, 17 β -hydroxy-5 α -estr-1-ene-3-one, and mixtures thereof.

5

[0021] The alkylcarbonate ester promoiety may be as described above. It may, for example, be selected from the group consisting of methyl carbonate, ethyl carbonate, propyl carbonate, isopropyl carbonate, butyl carbonate, isobutyl carbonate, t-butyl carbonate, valeryl carbonate, hexyl carbonate, heptyl carbonate, octyl carbonate, nonyl carbonate, decyl carbonate, undecyl carbonate, dodecyl carbonate, cyclopentylmethyl carbonate, cyclopentyl carbonate, cyclopentylpropyl carbonate, cyclohexylmethyl carbonate, cyclohexylpropyl carbonate, and mixtures thereof.

15

[0022] In accordance with this method aspect, the administered compound may comprise 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -alkylcarbonate (e.g., 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate), 17 β -hydroxyandrost-1,4-diene-3-one 17 β -alkylcarbonate (e.g., 17 β -hydroxyandrost-1,4-diene-3-one 17 β -ethylcarbonate), 17 β -hydroxy-5 α -estr-1-ene-3-one 17 β -alkylcarbonate (e.g., 17 β -hydroxy-5 α -estr-1-ene-3-one 17 β -ethylcarbonate), and/or 17 β -hydroxyestr-1,4-diene-3-one 17 β -alkylcarbonate (e.g., 17 β -hydroxyestr-1,4-diene-3-one 17 β -ethylcarbonate).

20

[0023] The compound to be administered may comprise diols, for example, such as 5 α -androst-1-ene-3,17 β -diol 17 β -alkylcarbonate (e.g., 5 α -androst-1-ene-3,17 β -diol 17 β -ethylcarbonate), 5 α -androst-1-ene-3,17 β -diol-3,17 β -di(alkylcarbonate) (e.g., 5 α -androst-1-ene-3,17 β -diol-3,17 β -di(ethylcarbonate), and mixtures thereof.

10051005001
10
[0024] The compound administration may comprise peroral administration, pernasal administration, transdermal administration, sublingual administration, and other means. Peroral administration is presently preferred.

5 [0025] As part of the method, the compound administration may comprise complexing the compound with an hydroxypropyl beta cyclodextrin, and/or with an hydroxypropyl gamma cyclodextrin.

[0026] Also as part of the compound administration according to this aspect of the invention, the compound preferably is administered once per day, although this is not limiting and other dosage regiments may be preferred depending upon the specific compound, the subject, etc. The compound may be administered in a dosage periodically for a maximum of two weeks, followed by at least two weeks of non-administration to permit recovery of natural parent androgen production in the subject.

15 [0027] The compound administration optionally but preferably comprises administering the compound in an amount ranging from 1.0 mg to 1 gram per day. More preferably, the compound administration comprises administering the compound in an amount ranging from 50 mg to 500 mg per day, and even more preferably in an amount ranging from 100 mg to 400 mg per day.

20 [0028] The compound administration optionally may further include applying an enteric coating to the compound prior to administering the compound.

DETAILED DESCRIPTION OF THE
PREFERRED EMBODIMENTS AND METHODS

[0029] Reference will now be made in detail to the presently preferred
embodiments and methods of the invention. It should be noted, however, that the
invention in its broader aspects is not limited to the specific details, representative
compositions and methods, and illustrative examples described in this section in
connection with the preferred embodiments and methods. The invention according
to its various aspects is particularly pointed out and distinctly claimed in the
attached claims read in view of this specification, and appropriate equivalents.

[0030] In accordance with one aspect of the invention, a compound is
provided for increasing the concentration of a parent androgen in a subject in vivo.
As noted above, the parent androgen has a skeletal structure including a 1 position
and a 17 position. This numbering corresponds to the ring identification and carbon
numbering system well known in the field of steroid chemistry, for example, as
adopted in the 1989 recommendations of the International Union of Pure and
Applied Chemistry ("IUPAC"). The parent androgen also has a 17 β -hydroxy group
comprising a 17 β -hydroxy oxygen appended to the 17 position and a 17 β -hydroxy
hydrogen appended to the 17 β -hydroxy oxygen.

[0031] The compound preferably but optionally is for treatment of a human
being to supplement or increase the concentration of the parent androgen in vivo.

This is not necessarily limiting, however, and veterinary applications also are possible in certain instances.

[0032] In accordance with this aspect of the invention, the compound comprises a substrate and a promoiety. The substrate has the skeletal structure of the parent androgen. It comprises a 1 position and a 17 position corresponding to the 1 and 17 positions respectively of the parent androgen. The substrate comprises a carbon-carbon double bond at the 1 position.

[0033] The skeletal structure of the parent androgen to which the substrate corresponds may take a number of forms. It may, for example, be selected from the group consisting of 5 α -androst-1-ene-3 α ,17 β -diol, 5 α -androst-1-ene-3 β ,17 β -diol, and mixtures thereof. It also may be selected from the group consisting of 5 α -estr-1-ene-3 α ,17 β -diol, 5 α -estr-1-ene-3 β ,17 β -diol, and mixtures thereof. In addition, it may be selected from the group consisting of 17 β -hydroxyandrost-1,4-diene-3-one 17 β -hydroxyestr-1,4-diene-3-one, 5 α ,17 β -hydroxyestr-1-ene-3-one, and mixtures thereof.

This skeletal structure may comprise, for example, 17 β -hydroxy-5 α -androst-1-ene-3-one.

[0034] The promoiety is appended to the 17 β -hydroxy oxygen of the substrate as a substitute for the 17 β -hydroxy hydrogen. It comprises and preferably consists of an alkylcarbonate ester. The alkyl group may be linear, branched, cyclical, etc. The promoiety preferably but optionally has an alkyl chain length (counting only the carbon atoms) of less than 12, and more preferably of less than

11. The alkylcarbonate ester may be selected from the group consisting of methyl carbonate, ethyl carbonate, propyl carbonate, isopropyl carbonate, butyl carbonate, isobutyl carbonate, t-butyl carbonate, valeryl carbonate, hexyl carbonate, heptyl carbonate, octyl carbonate, nonyl carbonate, decyl carbonate, undecyl carbonate, dodecyl carbonate, cyclopentyl methyl carbonate, cyclopentylpropyl carbonate, cyclohexyl methyl carbonate, cyclohexylpropyl carbonate, and mixtures thereof. Other alkyl carbonate esters, however, may be used. Alkyl carbonate esters having lower carbon chain lengths generally are preferred, although not universally so. Ethylcarbonate esters generally are even more preferred.

[0035] The compound according to this aspect of the invention in its presently preferred embodiments may comprise 17β -hydroxy- 5α -androst-1-ene-3-one 17β -alkylcarbonate, 17β -hydroxyandrost-1,4-diene-3-one 17β -alkylcarbonate. The ethyl carbonate esters, e.g., 17β -hydroxy- 5α -androst-1-ene-3-one 17β -ethylcarbonate, and 17β -hydroxyandrost-1,4-diene-3-one 17β -ethylcarbonate, again, would be preferred.

[0036] The compound also may comprise estranes, for example, such as 17β -hydroxy- 5α -estr-1-ene-3-one 17β -alkylcarbonate, 17β -hydroxyestr-1,4-diene-3-one 17β -alkylcarbonate, and their ethylcarbonate embodiments such as 17β -hydroxy- 5α -estr-1-ene-3-one 17β -ethylcarbonate. The compound also may comprise diols.

Illustrative examples would include 5α -androst-1-ene-3, 17β -diol 17β -alkylcarbonate (e.g., 5α -androst-1-ene-3, 17β -diol 17β -ethylcarbonate), and 5α -androst-1-ene-3, 17β -

diol-3,17 β -di(alkylcarbonate), (e.g., 5 α -androst-1-ene-3,17 β -diol-3,17 β -di(ethylcarbonate).

[0037] The principles of the invention according to this aspect also may be applied to dienes. Illustrative examples might include androst-1,4-diene-3 α ,17 β -diol 17 β -alkylcarbonate (e.g., androst-1,4-diene-3 α ,17 β -diol 17 β -ethylcarbonate), androst-1,4-diene-3 β ,17 β -diol 17 β -alkylcarbonate (e.g., androst-1,4-diene-3 β ,17 β -diol 17 β -ethylcarbonate), androst-1,4-diene-3 β ,17 β -diol-3-alkylcarbonate (e.g., androst-1,4-diene-3 β ,17 β -diol-3-ethylcarbonate), androst-1,4-diene-3 β ,17 β -diol-3,17 β -di(alkylcarbonate) (e.g., androst-1,4-diene-3 β ,17 β -diol-3,17 β -di(ethylcarbonate), and mixtures of these.

[0038] The compound may be contained or encapsulated by an enteric coating. The compound also may be administered with a carrier, which may comprise a solid carrier, a semi-solid carrier, or a liquid carrier. A preferred liquid carrier is an aqueous emulsion including fatty acid ethyl esters, polysorbate 60, lecithin, and cholesterol or an oil. Another preferred liquid carrier comprises water, glycerin, polysorbate, lecithin, sodium benzoate, ethylene diamine tetraacetic acid ("EDTA"), potassium sorbate, grapefruit seed extract, and vegetable gum.

[0039] The compound according to this aspect of the invention may and preferably does comprise combinations of compounds having the alkylcarbonate ester promoiety as described herein. As noted, in each instance in this document wherein a group is bonded at the 3 position (at the first or "A" carbon ring), the 3

position configuration may comprise or consist essentially of either the α configuration or the β , and preferably both. In addition, stereochemistry at the 5 position for compounds other than 4-ene compounds preferably comprise 5α .

[0040] In accordance with another aspect of the invention, a compound is provided for increasing the concentration of a parent androgen in a subject in vivo, wherein the parent androgen has a skeletal structure including a 4 position and a 17 position. The parent androgen also has a 17β -hydroxy group comprising a 17β -hydroxy oxygen appended to the 17 position and a 17β -hydroxy hydrogen appended to the 17β -hydroxy oxygen.

[0041] The compound according to this aspect of the invention comprises a substrate having the skeletal structure of the parent androgen comprising a 4 position and a 17 position corresponding to the 4 and 17 positions respectively of the parent androgen. The substrate comprises a carbon-carbon double bond at the 4 position. The compound further comprises a promoiety appended to the 17 position of the substrate as a substitute for the hydroxy hydrogen of the parent androgen. The promoiety comprises an alkylcarbonate ester.

[0042] The skeletal structure of the parent androgen to which the substrate corresponds may, for example, be selected from the group consisting of androst-4-ene- $3\alpha,17\beta$ -diol, androst-4-ene- $3\beta,17\beta$ -diol, and mixtures thereof. It also may be selected from the group consisting of estr-4-ene- $3\alpha,17\beta$ -diol, estr-4-ene- $3\beta,17\beta$ -diol, and mixtures thereof.

10043] The aspects of the alkylcarbonate ester promoiety as noted with
respect to the previous aspect of the invention apply here as well.

10044] Preferred but merely illustrative compounds according to this aspect
of the invention may comprise various forms of diols, for example, such as androst-
5 4-ene-3,17 β -diol 17 β -alkylcarbonate (e.g., androst-4-ene-3,17 β -diol 17 β -
ethylcarbonate), androst-4-ene-3,17 β -diol-3,17 β -di(alkylcarbonate) (androst-4-ene-
3,17 β -diol-3,17 β -di(ethylcarbonate)), estr-4-ene-3,17 β -diol 17 β -alkylcarbonate (e.g.,
estr-4-ene-3,17 β -diol 17 β -ethylcarbonate, estr-4-ene-3,17 β -diol-3,17 β -
di(alkylcarbonate) (e.g., estr-4-ene-3,17 β -diol-3,17 β -di(ethylcarbonate)).

10045] The compound according to this aspect of the invention also may
comprise a carrier, such as those forms noted above.

10046] In accordance with another aspect of the invention, a method is
provided for increasing the concentration of a parent androgen in a subject in vivo,
wherein the parent androgen has a skeletal structure including a 1 position and a
15 17 position and the parent androgen further has a 17 β -hydroxy group comprising a
17 β -hydroxy oxygen appended to the 17 position and a 17 β -hydroxy hydrogen
appended to the 17 β -hydroxy oxygen.

10047] The method according to this aspect of the invention comprises
administering to the subject a compound comprising a substrate and a promoiety.

20 The substrate has the skeletal structure of the parent androgen comprising a 1
position and a 17 position corresponding to the 1 and 17 positions respectively of the

parent androgen, and the substrate comprises a carbon-carbon double bond at the 1 position. The promoiety is appended to the 17 position of the substrate as a substitute for the 17 β -hydroxy hydrogen of the parent androgen, and comprises an alkylcarbonate ester.

5 **[0048]** In accordance with this aspect of the invention, the skeletal structure of the parent androgen to which the substrate corresponds may be selected from the group consisting of 5 α -androst-1-ene-3 α ,17 β -diol, 5 α -androst-1-ene-3 β ,17 β -diol, and mixtures thereof. It also may be selected from the group consisting of 5 α -estr-1-ene-3 α ,17 β -diol, 5 α -estr-1-ene-3 β ,17 β -diol, and mixtures thereof. The skeletal structure of the parent androgen to which the substrate corresponds also may comprise 17 β -hydroxy-5 α -androst-1-ene-3-one. It also may be selected from the group consisting of 17 β -hydroxyandrost-1,4-diene-3-one, 17 β -hydroxyestr-1,4-diene-3-one, 17 β -hydroxy-5 α -estr-1-ene-3-one, and mixtures thereof.

10
15 **[0049]** The alkylcarbonate ester promoiety may be as described above. It may, for example, be selected from the group consisting of methyl carbonate, ethyl carbonate, propyl carbonate, isopropyl carbonate, butyl carbonate, isobutyl carbonate, t-butyl carbonate, valeryl carbonate, hexyl carbonate, heptyl carbonate, octyl carbonate, nonyl carbonate, decyl carbonate, undecyl carbonate, dodecyl carbonate, cyclopentyl carbonate, cyclopentylmethyl carbonate, cyclopentylpropyl carbonate, cyclohexylmethyl carbonate, cyclohexylpropyl carbonate, and mixtures thereof.

20

[0050] In accordance with this method aspect, the administered compound may comprise 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -alkylcarbonate (e.g., 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate), 17 β -hydroxyandrost-1,4-diene-3-one 17 β -alkylcarbonate (e.g., 17 β -hydroxyandrost-1,4-diene-3-one 17 β -ethylcarbonate), 17 β -hydroxy-5 α -estr-1-ene-3-one 17 β -alkylcarbonate (e.g., 17 β -hydroxy-5 α -estr-1-ene-3-one 17 β -ethylcarbonate), and/or 17 β -hydroxyestr-1,4-diene-3-one 17 β -alkylcarbonate (e.g., 17 β -hydroxyestr-1,4-diene-3-one 17 β -ethylcarbonate).

[0051] The compound to be administered may comprise diols, for example, such as 5 α -androst-1-ene-3,17 β -diol 17 β -alkylcarbonate (e.g., 5 α -androst-1-ene-3,17 β -diol 17 β -ethylcarbonate), 5 α -androst-1-ene-3,17 β -diol 3,17 β -di(alkylcarbonate) (e.g., 5 α -androst-1-ene-3,17 β -diol 3,17 β -di(ethylcarbonate)).

[0052] In accordance with another aspect of the invention, a method is provided for increasing the concentration of a parent androgen in a subject in vivo, again, wherein the parent androgen has a skeletal structure including a 4 position and a 17 position and the parent androgen further has a 17 β -hydroxy group comprising a 17 β -hydroxy oxygen appended to the 17 position and a 17 β -hydroxy hydrogen appended to the 17 β -hydroxy oxygen.

[0053] The method according to this aspect of the invention comprises administering to the subject a compound comprising a substrate and a promoiety.

The substrate has the skeletal structure of the parent androgen comprising a 4 position and a 17 position corresponding to the 4 and 17 positions respectively of the

parent androgen, and the substrate comprises a carbon-carbon double bond at the 4 position. The promoiety is appended to the 17 position of the substrate as a substitute for the 17 β -hydroxy hydrogen of the parent androgen, and comprises an alkylcarbonate ester. The method further comprises converting the compound in vivo into the parent androgen. The subject again preferably but optionally is a human being, and the in vivo conversion thus correspondingly comprises converting the compound into the parent androgen in vivo within the human being.

[0054] The substrate has the skeletal structure of the parent androgen. The parent androgen may selected from the group consisting of androst-4-ene-3 α ,17 β -diol, androst-4-ene-3 β ,17 β -diol, and mixtures thereof. The parent androgen also may be selected from the group consisting of estr-4-ene-3 α ,17 β -diol, estr-4-ene-3 β ,17 β -diol, and mixtures thereof.

[0055] The alkylcarbonate ester promoiety may be as described above.

[0056] The compound also may comprise 4-ene diols, for example, such as androst-4-ene-3,17 β -diol 17 β -alkylcarbonate (e.g., androst-4-ene-3,17 β -diol 17 β -ethylcarbonate), androst-4-ene-3,17 β -diol 3,17 β -di(alkylcarbonate) (e.g., androst-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate)), estr-4-ene-3,17 β -diol 17 β -alkylcarbonate (e.g., estr-4-ene-3,17 β -diol 17 β -ethylcarbonate), estr-4-ene-3,17 β -diol 3,17 β -di(alkylcarbonate) (e.g., estr-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate)), and the like.

100570043001
[0057] In each of the aforementioned methods, the compound administration may comprise peroral administration, pernasal administration, transdermal administration, sublingual administration, and other means. The administration of the compound also may be by combinations of these techniques or approaches.

5 Peroral administration is presently preferred.

[0058] As part of the method, the compound administration may comprise complexing the compound with an hydroxypropyl beta cyclodextrin, and/or with an hydroxypropyl gamma cyclodextrin. This is particularly applicable if administered sublingually. The compound administration optionally may further include applying an enteric coating to the compound prior to administering the compound.

[0059] When administered orally or sublingually, the compound enters the gastrointestinal ("GI") tract, and ultimately the blood stream. Through more direct methods such as through pernasal, transdermal or intravenous injection, the compound enters directly into the blood stream. In each of the instances, the
15 compound may react to form the parent androgen or a prodrug of the parent androgen.

[0060] One limitation of known prodrugs of steriods is that, once they are tranformed into the parent steriod, they are broken down in the body, and particularly in the liver. This breakdown reduces in vivo concentration and
20 bioavailability of the steriod. In the presently preferred embodiments of the invention, the compounds are less prone to such breakdown in the body relative to

many known prodrug-type compounds. This in many instances is attributable to the alkylcarbonate ester promoiety, which makes the compound more resistant to hydrolysis and other reactions that inhibit or destroy them in the body. In vivo concentrations thus can be maintained more readily, and bioavailability of the parent androgen can be improved.

[0061] The compound preferably would be administered in amounts effective to supplement or increase the concentration of the parent androgen in vivo. The appropriate dosage therefore may take into account natural or otherwise expected variations in steroid concentration, such as the normal daily variations in natural steroid production and consumption, and such as normal variations in in vivo steroid concentrations over days or weeks.

[0062] According to a related aspect of the method, the compound may be administered using a dosage given periodically for a maximum of two weeks, followed by a period, for example, of at least two weeks, of non-administration to permit recovery of natural parent androgen production in the subject, for example, to the same baseline level possessed by the subject prior to use of the compound.

[0063] This can permit the compound to supplement or increase the concentration of the parent androgen in vivo for an effective period, and then terminate further dosages of the compound as its effectiveness attenuates.

[0064] The preferred dosage of the compound will depend upon the specific compound, the subject or class of subject to which it is to be administered, and other

factors commonly affecting dosage determinations for this type of composition. In general, the dosage should be such that a sufficient amount of the compound enters the system of the subject and supplements or increases the natural in vivo concentration of the parent androgen.

5 **[0065]** In accordance with presently preferred versions of the inventive method, the compound administration, particularly when applied to humans, comprises administering the compound in an amount ranging from 1.0 mg to 1 gram per day, more preferably in an amount ranging from 50 mg to 500 mg per day, and even more preferably in an amount ranging from 100 mg to 400 mg per day.

10 **[0066]** Also as part of the compound administration according to this aspect of the invention, the compound may be administered in a dosage periodically for a maximum of two weeks, followed by at least two weeks of non-administration to permit recovery of natural parent androgen production in the subject.

15 **[0067]** The compound administration also may comprise administering the compound only in morning-time.

20 **[0068]** Advantages of the preferred compounds and methods according to the invention flow from the advantageous or desirable effects of the parent androgen, as well as from the increased in vivo concentration and bioavailability of the prodrug and/or the corresponding parent androgen resulting from administration of the compound. Compounds such as the alkyl carbonate ester versions wherein the parent androgen skeletal structure corresponds to 5 α -androst-1-ene-3 α ,17 β -diol, 5 α -

androst-1-ene-3 β ,17 β -diol, and mixtures thereof, 5 α -estr-1-ene-3 α ,17 β -diol, 5 α -estr-1-ene-3 β ,17 β -diol, and mixtures thereof, 17 β -hydroxy-5 α -androst-1-ene-3-one, 17 β -hydroxyandrost-1,4-diene-3-one, 17 β -hydroxyestr-1,4-diene-3-one, 17 β -hydroxy-5 α -estr-1-ene-3-one, and mixtures thereof, androst-4-ene-3 α ,17 β -diol, androst-4-ene-3 β ,17 β -diol, and mixtures thereof, estr-4-ene-3 α ,17 β -diol, estr-4-ene-3 β ,17 β -diol, and mixtures thereof can provide pro-athletic and anabolic effects. In some instances, they can provide improved immune system efficacy, libido increase and/or mood enhancement as well.

[0069] Aspects of the alkylcarbonate ester as described above for the other aspects of the invention apply to the compound according to this aspect as well. A carrier such as those forms noted above also may be used.

EXAMPLE 1

[0070] To 10 ml dichloromethane was added 1.0 gram androst-4-ene-3,17-diol and 2.00 equivalents each of ethyl chloroformate and 2,6-lutidine. This was then stirred overnight, diluted with an additional 60 ml of dichloromethane, and filtered. The filtrate was washed twice in a separatory funnel with acidic water, once with neutral water, then dried over sodium sulfate and placed in a freezer at -10 C for two hours. The resulting white colorless crystals were recovered by filtration, yielding 0.47 grams of androst-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) ester.

[0071] Pyridine or triethylamine may be used as alternate bases. Alternate solvents include acetone, acetonitrile, and pyridine.

EXAMPLE 2

[0072] A compound according to this aspect of the invention may be made from what in fact is comprised of a combination of compounds according to the invention. About 100 mg of a first component of androst-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) and about 200 mg of a second component of 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate ester are added to a liquid carrier component comprising 2,876 mg of purified water, 2,443 mg of glycerin, 1,847 mg of propylene glycol, 459 mg of polysorbate, 396 mg of Lecithin, 202 mg of EDTA. 185 mg of vegetable gum, 566 mg of natural flavor blend, 96 mg of sodium benzoate, 96 mg of potassium sorbate, and 96 mg of calcium propionate. The components are stirred to achieve dissolution and thus form the resultant solution. This specific example yields approximately 9,620 mg or about 9 ml of the solution. This solution can be taken by a human subject in daily doses of about 9-18 ml.

EXAMPLE 3

[0073] About 150 mg of androst-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) and about 150 mg of 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate ester are added to a liquid carrier component comprising 2,876 mg of purified water, 2,443 mg of glycerin, 1,847 mg of propylene glycol, 459 mg of polysorbate, 396 mg of Lecithin, 202 mg of EDTA. 185 mg of vegetable gum, 566 mg of natural flavor

blend, 96 mg of sodium benzoate, 96 mg of potassium sorbate, and 96 mg of calcium propionate. The components are stirred to achieve dissolution and thus form the resultant solution. This specific example yields approximately 9,620 mg or about 9 ml of the solution. This solution can be taken by a human subject in daily doses of about 9-18 ml.

EXAMPLE 4

[0074] About 200 mg of androst-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) and about 100 mg of 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate ester are added to a liquid carrier component comprising 2,876 mg of purified water, 2,443 mg of glycerin, 1,847 mg of propylene glycol, 459 mg of polysorbate, 396 mg of Lecithin, 202 mg of EDTA. 185 mg of vegetable gum, 566 mg of natural flavor blend, 96 mg of sodium benzoate, 96 mg of potassium sorbate, and 96 mg of calcium propionate. The components are stirred to achieve dissolution and thus form the resultant solution. This specific example yields approximately 9,620 mg or about 9 ml of the solution. This solution can be taken by a human subject in daily doses of about 9-18 ml.

EXAMPLE 5

[0075] About 150 mg of androst-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) and about 150 mg of 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate ester are added to a liquid carrier component comprising 2,150 mg of purified water, 2,800 mg of glycerin, 2,216 mg of propylene glycol, 459 mg of polysorbate, 396 mg of

Lecithin, 202 mg of EDTA. 185 mg of vegetable gum, 566 mg of natural flavor blend, 96 mg of sodium benzoate, 96 mg of potassium sorbate, and 96 mg of calcium propionate. The components are stirred to achieve dissolution. This yields approximately 9,620 mg or about 9 ml of the solution. This solution can be taken by
5 a human subject in daily doses of about 9-18 ml.

EXAMPLE 6

[0076] About 150 mg of androst-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) and about 150 mg of 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate ester are added to a liquid carrier component comprising 3,550 mg of purified water, 2,132 mg of glycerin, 1,484 mg of propylene glycol, 459 mg of polysorbate, 396 mg of
10 Lecithin, 202 mg of EDTA. 185 mg of vegetable gum, 566 mg of natural flavor blend, 96 mg of sodium benzoate, 96 mg of potassium sorbate, and 96 mg of calcium propionate. The components are stirred to achieve dissolution. This yields approximately 9,620 mg or about 9 ml of the solution. This solution can be taken by
15 a human subject in daily doses of about 9-18 ml.

EXAMPLE 7

[0077] About 100 mg of a first component of estr-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) and about 200 mg of a second component of 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate ester are added to a liquid carrier
20 component comprising 2,876 mg of purified water, 2,443 mg of glycerin, 1,847 mg of propylene glycol, 459 mg of polysorbate, 396 mg of Lecithin, 202 mg of EDTA. 185

mg of vegetable gum, 566 mg of natural flavor blend, 96 mg of sodium benzoate, 96 mg of potassium sorbate, and 96 mg of calcium propionate. The components are stirred to achieve dissolution and thus form the resultant solution. This specific example yields approximately 9,620 mg or about 9 ml of the solution. This solution can be taken by a human subject in daily doses of about 9-18 ml.

EXAMPLE 8

[0078] About 150 mg of estr-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) and about 150 mg of 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate ester are added to a liquid carrier component comprising 2,876 mg of purified water, 2,443 mg of glycerin, 1,847 mg of propylene glycol, 459 mg of polysorbate, 396 mg of Lecithin, 202 mg of EDTA. 185 mg of vegetable gum, 566 mg of natural flavor blend, 96 mg of sodium benzoate, 96 mg of potassium sorbate, and 96 mg of calcium propionate. The components are stirred to achieve dissolution and thus form the resultant solution. This yields approximately 9,620 mg or about 9 ml of the solution, which can be taken by a human subject in daily doses of about 9-18 ml.

EXAMPLE 9

[0079] About 200 mg of estr-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) and about 100 mg of 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate ester are added to a liquid carrier component comprising 2,876 mg of purified water, 2,443 mg of glycerin, 1,847 mg of propylene glycol, 459 mg of polysorbate, 396 mg of Lecithin, 202 mg of EDTA. 185 mg of vegetable gum, 566 mg of natural flavor

blend, 96 mg of sodium benzoate, 96 mg of potassium sorbate, and 96 mg of calcium propionate. The components are stirred to achieve dissolution and thus form the resultant solution. This specific example yields approximately 9,620 mg or about 9 ml of the solution. This can be taken by a human subject in daily doses of about 9-18 ml.

EXAMPLE 10

[0080] About 150 mg of estr-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) and about 150 mg of 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate ester are added to a liquid carrier component comprising 2,150 mg of purified water, 2,800 mg of glycerin, 2,216 mg of propylene glycol, 459 mg of polysorbate, 396 mg of Lecithin, 202 mg of EDTA. 185 mg of vegetable gum, 566 mg of natural flavor blend, 96 mg of sodium benzoate, 96 mg of potassium sorbate, and 96 mg of calcium propionate. The components are stirred to achieve dissolution. This yields approximately 9,620 mg or about 9 ml of the solution. This solution can be taken by a human subject in daily doses of about 9-18 ml.

EXAMPLE 11

[0081] About 150 mg of estr-4-ene-3,17 β -diol 3,17 β -di(ethylcarbonate) and about 150 mg of 17 β -hydroxy-5 α -androst-1-ene-3-one 17 β -ethylcarbonate ester are added to a liquid carrier component comprising 3,550 mg of purified water, 2,132 mg of glycerin, 1,484 mg of propylene glycol, 459 mg of polysorbate, 396 mg of Lecithin, 202 mg of EDTA. 185 mg of vegetable gum, 566 mg of natural flavor

blend, 96 mg of sodium benzoate, 96 mg of potassium sorbate, and 96 mg of calcium propionate. The components are stirred to achieve dissolution. This yields approximately 9,620 mg or about 9 ml of the solution. This solution can be taken by a human subject in daily doses of about 9-18 ml.

5 **[0082]** It has been noted that the compound may, and in many instances preferably does, comprise or consist of combinations of the compounds described herein and/or recited in the claims. The substrate is described in some instances herein as comprising a double bond at a particular position. Preferably in these instances the substrate includes a double bond in the first carbon ring (A) only at the designated position. The promoiety is described herein above as comprising an alkyl carbonate. Optionally and preferably, the promoiety consists of an alkyl carbonate, such as those identified and/or described herein.

10 **[0083]** Additional advantages and modifications will readily occur to those skilled in the art. Therefore, the invention in its broader aspects is not limited to the specific details, representative devices and methods, and illustrative examples shown and described. Accordingly, departures may be made from such details without departing from the spirit or scope of the general inventive concept as defined by the appended claims and their equivalents.